

Instantons and affine Lie algebras*

Hiraku Nakajima

Department of Mathematical Sciences, University of Tokyo,
3-8-1 Komaba, Meguro-ku, Tokyo 153, Japan

September, 1995

Abstract

Various constructions of the affine Lie algebra action on the moduli space of instantons on 4-manifolds are discussed. The analogy between the local-global principle and the role of mass is also explained. The detailed proofs are given in separated papers [16, 17].

1 Introduction

Vafa and Witten [20] introduced topological invariants¹ for 4-manifolds using $N = 4$ topological supersymmetric Yang-Mills theory. Then the S -duality conjecture implies that the generating function of those invariants is a modular form of certain weight, where the summation runs over all $SU(2)$ or $SO(3)$ -principal bundles of any topological types. (In general, it has the modular invariance only for $\Gamma_0(4)$, a subgroup of $SL(2, \mathbb{Z})$.) For some 4-manifolds, they identify those invariants with the Euler numbers of the instanton moduli spaces. Then they can check the modular invariance for various 4-manifolds, using mathematical results, Yoshioka's formulae [23] for $\mathbb{P}^2$ and the blow up, Göttsche and Huybrechts's result [5] for the K3 surface, and the author's result for ALE spaces.

*to appear in Proceedings of Trieste Conference on S-duality and Mirror Symmetry

¹The author does not know how to define their invariants in a mathematically rigorous way. The difficulty lies in the lack of the compactness of relevant moduli spaces.

On the other hand, the author's motivation of the study [14, 15, 16] of the homology groups of the instanton moduli spaces on ALE spaces is totally different. The author's motivation was trying to understand Ringel [19] and Lusztig's [11, 12] constructions of the lower triangular part $\mathbf{U}_q^-$ of the quantized enveloping algebra. They used the moduli spaces of representations of quivers, and their cotangent bundle² can be identified with the instanton moduli spaces on ALE spaces, via the ADHM description [10]. The author showed that the generating function of the Euler numbers of the instanton moduli spaces on ALE spaces becomes the character of the affine Lie algebra, which has been known to have modular transformation property by Kac-Peterson (see [8]).

The definition of the affine Lie algebra representation on the homology group of the instanton moduli spaces is very geometric, and seems to be generalized, at least, to projective surfaces. The results of similar direction are announced recently by Ginzburg-Kapranov-Vasserot [3] and Grojnowski [6]. Unfortunately, our construction depends heavily on the complex structure of the base manifold. It is a challenging problem to generalize the construction to more general 4-manifolds. One may need to reformulate the homology group of the moduli spaces, etc.....

Acknowledgements

The author's understanding of the analogy between the local-global principle and the theory of mass came from lectures given by the seminar on Seiberg-Witten theory organized by K. Ueno. He would like to speakers, especially S.-K. Yang. He also thank to G. Moore and K. Yoshioka for valuable discussions.

2 The Hilbert scheme of points and the Heisenberg algebra: Twist around points

In this section, we study the relationship between the Hilbert scheme of points and the Heisenberg algebra. The reasons why we study the Hilbert

²They are not cotangent bundles rigorously. The situation is very much similar to the relation between the moduli space of vector bundles over a curve and Hitchin's moduli space of Higgs bundles.

scheme are (a) it is a toy model for moduli spaces of instantons, (b) it appears in the boundary of the compactification of the instanton moduli spaces over projective surfaces, and (c) its homology group is isomorphic to that of a moduli space for some special cases [5].

We explain the reason (b) a little bit more. Since the instanton moduli spaces are usually noncompact, one must compactify them to consider their Euler numbers. When the base manifold is a projective surface, the results of Donaldson and Uhlenbeck-Yau enable us to identify the instanton moduli space with the moduli space of μ -stable holomorphic vector bundles (Hitchin-Kobayashi correspondence). Then the one of the most natural compactifications seems to be Gieseker-Maruyama's compactifications $\overline{\mathfrak{M}}$ ³, namely moduli spaces of semi-stable torsion free sheaves.

If $\mathcal{E}$ is a torsion free sheaf which is not locally free, its double dual $\mathcal{E}^{\vee\vee}$ is a locally free sheaf and we have an exact sequence

$$0 \rightarrow \mathcal{E} \rightarrow \mathcal{E}^{\vee\vee} \rightarrow \mathcal{E}^{\vee\vee}/\mathcal{E} \rightarrow 0.$$

Thus $\mathcal{E}$ can be determined by (a) $\mathcal{E}^{\vee\vee}$ and (b) $\mathcal{E}^{\vee\vee} \rightarrow \mathcal{E}^{\vee\vee}/\mathcal{E}$. The double dual $\mathcal{E}^{\vee\vee}$ is contained in the interior of $\overline{\mathfrak{M}}$, but in the different component with lower second Chern number. Thus it is natural to expect that those studies can be decomposed into two parts, the interior (a) and the quotient map (b). And the variety of the quotient map (b), which depends only on the rank of $\mathcal{E}$ and the length of $\mathcal{E}^{\vee\vee}/\mathcal{E}$, looks very much like the Hilbert scheme of points. In fact, the Hilbert scheme is the special case $\mathcal{E}^{\vee\vee} = \mathcal{O}$. The Betti numbers of the variety was computed by Yoshioka [23, 0.4].

Let X be a projective surface defined over $\mathbb{C}$. Let $X^{[n]}$ be the component of the Hilbert scheme of X parameterizing the ideals of $\mathcal{O}_X$ of colength n . It is smooth and irreducible. Let $S^n X$ denotes the n -th symmetric product of X . It parameterizes formal linear combinations $\sum n_i [x_i]$ of points x_i in X with coefficients $n_i \in \mathbb{Z}_{>0}$ with $\sum n_i = n$. There is a canonical morphism $\pi: X^{[n]} \rightarrow S^n X$ defined by

$$\pi(\mathcal{J}) = \sum_{x \in X} \text{length}(\mathcal{O}_X/\mathcal{J})_x [x].$$

It is known that π is a resolution of singularities.

³The Gieseker-Maruyama's compactifications are not smooth in general. In fact, Vafa-Witten's formula for the K3 surface gives the fractional Euler number. This may be the contribution of the singularities.

Göttsche [4] computed the generating function of the Poincaré polynomials

$$\begin{aligned} & \sum_{n=0}^{\infty} q^n P_t(X^{[n]}) \\ &= \prod_{m=1}^{\infty} \prod_{i=0}^4 (1 - (-t)^{2m-2+i} q^m)^{(-1)^{i+1} b_i(X)}, \end{aligned}$$

where $b_i(X)$ is the Betti number of X . It was shown that the Euler number of $X^{[n]}$ is equal to the orbifold Euler number of $S^n X$ by Hirzebruch-Hofer [7]. It was also pointed out by Vafa and Witten that this is equal to the character of the Fock space.

We shall construct the representation of the Heisenberg and Clifford algebras in a geometric way. The key point is to introduce appropriate ‘‘Hecke correspondence’’ which give the generators of the Heisenberg/Clifford algebra.

Take a basis of $H_*(X)$ and assume that each element is represented by a (real) compact submanifold C^a . (a runs over $1, 2, \dots, \dim H_*(X)$.) Take a dual basis for $H_*(X)$, and assume that each element is also represented by a submanifold D^a . (Those assumptions are only for the brevity. The modification to the case of cycles is clear.) For each $a = 1, 2, \dots, \dim H_*(X)$, $n = 1, 2, \dots$ and $i = 1, 2, \dots$, we introduce cycles of products of the Hilbert schemes by

$$\begin{aligned} E_i^a(n) &= \{ (\mathcal{J}_1, \mathcal{J}_2) \in X^{[n-i]} \times X^{[n]} \mid \mathcal{J}_1 \supset \mathcal{J}_2 \\ &\quad \text{and } \text{Supp}(\mathcal{J}_1/\mathcal{J}_2) = \{p\} \text{ for some } p \in D^a \}, \\ F_i^a(n) &= \{ (\mathcal{J}_1, \mathcal{J}_2) \in X^{[n+i]} \times X^{[n]} \mid \mathcal{J}_1 \subset \mathcal{J}_2 \\ &\quad \text{and } \text{Supp}(\mathcal{J}_2/\mathcal{J}_1) = \{p\} \text{ for some } p \in C^a \}. \end{aligned}$$

Then we define an endomorphism $H_*(X^{[n]}) \rightarrow H_*(X^{[n-i]})$ by

$$c \mapsto (p_1)_*([E_i^a(n)] \cap p_2^*c),$$

where p_1, p_2 are projections of the first and second factor of $X^{[n-i]} \times X^{[n]}$ and $p_2^*c = [X^{[n-i]}] \times c$ and $(p_1)_*$ is a push-forward. Similarly, we have an endomorphism $H_*(X^{[n]}) \rightarrow H_*(X^{[n+i]})$ using $F_i^a(n)$. Collecting the operators with respect to n , we have operators $[E_i^a]$, $[F_i^a]$ acting on the direct sum $\bigoplus_n H_*(X^{[n]})$. Then

Theorem 1 . *The following relations hold as operators on $\bigoplus_n H_*(X^{[n]})$.*

$$\begin{aligned} [E_i^a][E_j^b] &= (-1)^{\dim D^a \dim D^b} [E_j^b][E_i^a] \\ [F_i^a][F_j^b] &= (-1)^{\dim C^a \dim C^b} [F_j^b][F_i^a] \\ [E_i^a][F_j^b] &= (-1)^{\dim D^a \dim C^b} [F_j^b][E_i^a] \\ &\quad + \delta_{ab} \delta_{ij} c_i \text{Id}, \end{aligned}$$

where c_i is a nonzero integer depending only on i (independent of X).

In particular, for each fixed a , $[E_i^a]$, $[F_i^a]$ ($i = 1, \dots$) define the action of the Heisenberg or Clifford algebra according to the parity of $\dim C^a$.

Moreover, comparing with Göttsche's formula, we can conclude our representation is irreducible.

The definition of the correspondence E_i^a , F_i^a can be naturally generalized to the case of moduli spaces of torsion free sheaves (see [6]). However, the author has no idea to generalize to more general 4-manifolds.

3 Elementary transformation: Twist along embedded submanifolds

The operator of the previous section twists sheaves around points. There is another kind of operator which twists sheaves along an embedded 2-dimensional submanifold. This operation is called the elementary transformation in the literature.

Suppose C is a holomorphic curve embedded in a projective surface X . Let i denote the inclusion map. Let $\mathfrak{M}$ be the moduli space of $U(r)$ -instantons, namely Einstein-Hermitian connections. We identify it with the moduli space of holomorphic vector bundles over X by the Hitchin-Kobayashi correspondence. It decomposes by the first and second Chern classes. For each integer d , we also consider the following moduli space $\mathfrak{P}$ of parabolic bundles $(\mathcal{E}_1, \mathcal{E}_2, \varphi)$ where $\mathcal{E}_i$ is a holomorphic vector bundle over X , and $\varphi: \mathcal{E}_1 \rightarrow \mathcal{E}_2$ is an injection which is an isomorphism outside C . In order to define the moduli space, we need to introduce the notion of the stability to parabolic bundles (see [13]), for this we need to choose an ample line bundle L or the Kähler metric which is a curvature of L . Moreover, it is necessary

to consider the Gieseker-Maruyama compactification of moduli spaces, as explained in the previous section. But we do not go in detail.

There is a morphism $f: \mathfrak{P} \rightarrow \mathfrak{M} \times \mathfrak{M}$. Then we can define two operators on the homology group exactly as in the previous section:

$$\begin{aligned} c &\mapsto (p_1)_*(f_*[\mathfrak{P}] \cap p_2^*c), \\ c &\mapsto (p_2)_*(p_1^*c \cap f_*[\mathfrak{P}]) \end{aligned}$$

Note that the first and second Chern classes are not preserved this operator. Strictly speaking, we do not have a globally defined morphism since the stability conditions for the parabolic bundles and their underlying vector bundles are not equivalent in general. But it is enough for our purpose to have $f_*[\mathfrak{P}]$ as an element of homology group of $\mathfrak{M} \times \mathfrak{M}$. For example, f could be a meromorphic map.

Since we do not know what is the right setting for general projective surfaces⁴, we focus on particular examples, namely ALE spaces. The ALE spaces are the minimal resolution of simple singularities $\mathbb{C}^2/\Gamma$, where Γ is a finite subgroup of $SU(2)$. The second homology group H_2 of the ALE space is spanned by the irreducible components $\Sigma_1, \dots, \Sigma_n$ of the exceptional set, which are the projective line. The intersection matrix is the negative of the Cartan matrix of type ADE. The classification of simple singularities are given by the Dynkin graphs in this way. In particular, there is a bijection between simple singularities and simple Lie algebra of type ADE. The rank is equal to the number of the number of the irreducible components, namely n . By the work of Kronheimer, it is known that they have hyper-Kähler metrics. There are a variant of the ADHM description, which identifies the framed moduli spaces of instantons, or more precisely, torsion-free sheaves with the cotangent bundles of the moduli space of representations of quivers of affine Dynkin graphs.

Since we shall work on non-compact spaces, we have extra discrete parameters which parameterizes the boundary condition. We consider instantons which converge to a flat connection at the end of the ALE space X . The flat connection on the end can be classified by its monodromy, namely a representation ρ of the finite group Γ . Let $\rho_0, \rho_1, \dots, \rho_n$ be the irreducible

⁴If one could define the Hecke operators using Kronheimer-Mrowka's singular anti-self-dual connections, they might be the right setting.

representations of Γ with ρ_0 the trivial representation. By the McKay correspondence, there is a bijection between the vertices of the affine Dynkin graph and the irreducible representations (see [15] for more detail). The monodromy representation ρ is decomposed as $\rho = \bigoplus_{k=0}^n \rho_k^{\oplus w_k}$, where w_k is the multiplicity. This datum will be preserved under the Hecke operator.

Corresponding to each irreducible component Σ_k , we take a component of the moduli space of parabolic bundles where $\mathcal{E}_2/\varphi(\mathcal{E}_1)$ is rank 1 and degree -1 . We then define operators e_k and f_k on the homology group of the moduli space as above. We also have an operator $\alpha_k^\vee$ which is the multiplication by $-\langle c_1, [\Sigma_k] \rangle$ on the homology class belonging to the component with the first Chern class c_1 .

For $k = 0$, we can define similar operators e_0, f_0 by replacing $\mathcal{O}_{\Sigma_k}(-1)$ by a sheaf $\mathcal{O}_{\bigcup \Sigma_k}$. The operator $\alpha_0^\vee$ is defined so that

$$\sum_{k=0}^n \dim \rho_k \alpha_k^\vee = \text{rank } E \text{ Id}.$$

Finally define the operator d to detect the instanton number. Namely the multiplication by

$$- \int_X \text{ch}(\mathcal{E}).$$

on the homology group of the each component of the moduli spaces.

Theorem 2 . Operators $\alpha_k^\vee, e_k, f_k$ ($k = 0, \dots, n$), d satisfy the relation of the affine Lie algebra corresponding to the extended Dynkin graph. Moreover, the representation on $H_*(\mathfrak{M})$ is integrable.

The irreducible decomposition of the representation is complicated, but we have one irreducible factor whose geometric meaning is clear.

Theorem 3 . If we take the middle degree part of the homology group $H_*(\mathfrak{M})$ (since the dimension of $\mathfrak{M}$ are changing on components, the middle degree also changes), it is preserved by the affine Lie algebra action. Moreover, it is the integrable highest weight representation with the highest weight vector ${}^t(w_0, \dots, w_n)$. The level is equal to the rank of the vector bundle.

The highest weight vector lies in the particularly chosen moduli space which consists of a single point.

4 Local-Global Principle and the Mass

Historically the Hecke operators were originally introduced in the theory of modular forms. There are also analogous operators in the theory of the moduli spaces vector bundles over curves [18], which are used in the geometric Langlands program. Our operators can be considered as natural complex 2-dimensional analogue of these operators.

The importance of the Hecke operators comes from the fact that they lie in the heart of the “local-global principle”. We shall explain it only very briefly. The interested reader should consult to good literatures about the Hecke operators and the modular forms (see e.g., [2]).

The local-global principle roughly says that a global problem could be studied as a collection of local problems. The basic example is the Hasse principle: A quadratic form with integer coefficients has a nontrivial integer solution if and only if it has real solution and a p -adic solution. In this case the global problem is to find an integer solution and the local problems are find solutions in $\mathbb{R}$ and $\mathbb{Q}_p$. Thus the base manifold, which is the parameter space of the local places, is the set of prime numbers plus infinity $\mathbb{R}$.

The theory of modular forms are also examples of the global-local principle. Consider the space of modular forms of weight k , which can be considered as functions of lattices in $\mathbb{C}$ with homogeneous degree $-k$, i.e., $F(\lambda L) = \lambda^{-k} F(L)$. Then for each prime number p , we define the Hecke operator $T(p)$ by

$$(T(p)F)(L) = p^{k-1} \sum_{[L':L]=p} F(L'),$$

where the summation runs over the set of sublattices of L with index p . These operators commute each other. If a modular form is a simultaneous eigenfunction, its L -function has an Euler product expansion. The analogy between the modular forms and 4-dimensional gauge theory our theory are given in the table.

In physics, there is a good explanation why the local-global principle holds in some topological field theories⁵. In these theories, topological invariants, like Donaldson’s invariants, are expressed as correlation functions. It is not by no means obvious that the results are topological invariants, since one

⁵Topological field theories of cohomological type according to the terminology in [1]

prime numbers	points in a 4-manifold and submanifolds C
the space of modular forms	the homology group of moduli spaces
Hecke operators	our Hecke operators

needs to introduce a Riemannian metric for the definition of the Lagrangian. However, by a clever choice of the Lagrangian, the resulted correlation functions are independent of the choice of the metric. The mechanism is just like the fact that the euler class, which is defined as the pfaffian of the curvature, is a topological invariant. Thus one can take a family of Riemannian metrics $g_t = tg$ with $t > 0$, and consider the limiting behaviour for $t \rightarrow \infty$. In the limit, the distance of two different points goes to infinity. Hence if the “mass” of all particles is not zero, there are no interaction between two points. Then one can compute the correlation functions by integrating local contributions over the base manifold. In this sense, the local-global principle holds in this theory.

In $N = 1$ topological supersymmetric Yang-Mills theory, it is believed that all particles have non-trivial mass. However in $N = 2$ topological supersymmetric Yang-Mills theory, which is relevant to Donaldson’s invariants, it is no longer true. Hence there may be massless particles which make interaction even when the distance between two points are very large. Anyway, if all particles would have non-trivial mass, Donaldson’s invariants would depend only on homology classes of the underlying manifolds. On a Kähler manifold X with a non-trivial holomorphic 2-form ω , Witten [21] used a perturbation from the $N = 2$ theory to the $N = 1$ theory adding a term depending on ω . The remarkable observation was that there remain particles which have zero mass where ω vanishes. Unless the manifold X is a K3 surface or a torus, ω vanishes along a divisor C . Thus the local-global principle holds in the $N = 2$ theory with only one modification; there are non-trivial contribution from C . In other words, the 2-dimensional submanifold C cannot be divided any more, and should be considered as a point. Similarly, in [20], again for the same class of Kähler manifolds, the $N = 4$ theory was perturbed to the $N = 1$ theory, and the correlation function was calculated in a similar way. And finally, Witten conjectured [22] that the Kronheimer-Mrowka’s basic classes [9] coincide with homology classes whose Seiberg-Witten invariants

are nonzero. It means that the local-global principle fails exactly along basic classes.

Now it becomes clear why we must introduce two kinds of Hecke correspondences, twist along a point and twist along a 2-dimensional submanifold. For the Hilbert scheme on a projective surface, it is apparent that the local-global principle holds without the introduction of C . This should be the basic reason why the homology group of the Hilbert scheme is generated only by the first kind of the Hecke correspondence. For higher rank case, we might need the second type of the Hecke correspondence in order to get all homology classes in the moduli spaces. However, the relation between two kinds of Hecke operators is not clear, at this moment. Moreover, there might be other types of correspondences which is useful to describe the homology group of the moduli space. For example, Göttsche and Huybrechts [5] used an interesting correspondence in order to relate moduli spaces of rank 2 bundles and Hilbert schemes.

Finally, we would like to point out the difference between our situation and the classical one (i.e., the Hecke operators on modular forms). The first kind of the Hecke operator is independent of the choice of the representative C^a of the homology class. This is because we are studying the homology group of the moduli space. It is not clear that the second kind of the Hecke operator depends only on the homology class of C . But it depends only on the rational equivalence class.

References

- [1] S. Cordes, G. Moore, and S. Ramgoolam, *Lectures on 2D Yang-Mills theory, equivariant cohomology and topological field theories*, hep-th/9411210.
- [2] S. Gelbart, *An elementary introduction of the Langlands program*, Bull. of the AMS, **10** (1984), 177–219.
- [3] V. Ginzburg, M. Kapranov, and E. Vasserot, *Langlands reciprocity for algebraic surfaces*, Int. Math. Res. Notes **2** (1995), 147–160.
- [4] L. Göttsche, *The Betti numbers of the Hilbert scheme of points on a smooth projective surface*, Math. Ann. **286** (1990), 193–207.

- [5] L. Göttsche and D. Huybrechts, *Hodge numbers of moduli spaces of stable bundles on K3 surfaces*, alg-geom/9408001, preprint.
- [6] I. Grojnowski, *Instantons and affine algebras I: the Hilbert scheme and vertex operators*, alg-geom/9506020, preprint.
- [7] F. Hirzebruch and T. Höfer, *On the Euler number of an orbifold*, Math. Ann. **286** (1990), 255–260.
- [8] V.G. Kac, *Infinite dimensional Lie algebras* (3rd Ed.), Cambridge Univ. Press 1990.
- [9] P.B. Kronheimer and T. Mrowka, *Embedded surfaces and the structure of Donaldson’s polynomial invariants*, J. Diff. Geom. (1995), 573–734.
- [10] P.B. Kronheimer and H. Nakajima, *Yang-Mills instantons on ALE gravitational instantons*, Math. Ann. **288** (1990), 263–307.
- [11] G. Lusztig, *Canonical bases arising from quantized enveloping algebras*, J. Amer. Math. Soc. **3** (1990), 447–498.
- [12] ———, *Quivers, perverse sheaves, and quantized enveloping algebras*, J. Amer. Math. Soc. **4** (1991), 365–421.
- [13] M. Maruyama and K. Yokogawa, *Moduli of parabolic stable sheaves*, Math. Ann. **293** (1992), 77–99.
- [14] H. Nakajima, *Instantons on ALE spaces, quiver varieties, and Kac-Moody algebras*, Duke Math. **76** (1994), 365–416.
- [15] ———, *Gauge theory on resolution of simple singularities and simple Lie algebras*, Inter. Math. Res. Notices (1994), 61 – 74.
- [16] ———, *Quiver varieties and Kac-Moody algebras*, preprint.
- [17] ———, *Heisenberg algebra and Hilbert schemes of points of projective surfaces*, alg-geom/9507012, preprint.
- [18] M.S. Narasimhan and S. Ramanan, *Deformation of the moduli space of vector bundles over an algebraic curve*, Ann. of Math. **101** (1975), 391–417.

- [19] C.M. Ringel *Hall algebras and quantum groups*, Invent. Math. **101** (1990), 583–592.
- [20] C. Vafa and E. Witten, *A strong coupling test of S-duality*, Nucl. Phys. **431** (1994), 3–77.
- [21] E. Witten, *Supersymmetric Yang-Mills theory on a four-manifold*, J. Math. Phys. **35** (1994), 5101–5135.
- [22] ———, *Monopoles and four-manifolds*, Int. Math. Res. Notes **1** (1995), 769–796.
- [23] K. Yoshioka, *The Betti numbers of the moduli spaces of stable sheaves of rank 2 on $\mathbb{P}^2$* , J. reine angew. Math. **453** (1994), 193–220.